

<https://doi.org/10.52889/1684-9280-2021-3-59-4-8>

УДК 617.3; 616-089.23; 616-001; 615.477.2

МРПТИ: 76.29.41

Review article

Epidemiological Medical-Social Aspects of Developmental Dysplastic Coxarthrosis

Durdana Oktyabrova ¹, Kairat Ashimov ², Timur Baidalin ³, Bekzhan Suleimenov ⁴, Askar Beknazarov ⁵

¹ Doctor-resident of the Department of orthopedics No.3, National Scientific Center of Traumatology and Orthopaedics named after Academician Batpenov N.D., PhD student of the Astana medical university, Nur-Sultan, Kazakstan.

E-mail: oktyabrovdana@gmail.com

² Head of the Department orthopedics No.3, National Scientific Center of Traumatology and Orthopaedics named after Academician Batpenov N.D., Nur-Sultan, Kazakstan. E-mail: kairat_ashimov@mail.ru

³ Head of the Department orthopedics No.7, National Scientific Center of Traumatology and Orthopaedics named after Academician Batpenov N.D., Nur-Sultan, Kazakstan. E-mail: t.baidalin@gmail.com

⁴ Traumatologist-orthopedist of the Department of orthopedics No.3, National Scientific Center of Traumatology and Orthopaedics named after Academician Batpenov N.D., Nur-Sultan, Kazakstan. E-mail: bekjan-suleimen@mail.ru

⁵ Traumatologist-orthopedist of the Department of orthopedics No.3, National Scientific Center of Traumatology and Orthopaedics named after Academician Batpenov N.D., Nur-Sultan, Kazakstan. E-mail: askar_big_man@mail.ru

Abstract

Dysplastic coxarthrosis incidence rates in the world differ due to geographic location and racial groups variety. Rates are higher for female, than they are for male. The number of patients with dysplastic coxarthrosis who are waiting for conducting total hip arthroplasty is increasing every day.

There is a necessity of detailed research of prevalence, as well as conducting medical and social evaluation of consequences among patients with dysplastic coxarthrosis in Kazakhstan.

Key words: developmental dysplasia of the hip, dysplastic coxarthrosis, life quality, disablement, young age.

Corresponding author: Durdana Oktyabrova, Doctor-resident of the National Scientific Center of Traumatology and Orthopedics named after Academician N.D. Batpenov, Nur-Sultan, Kazakhstan.

Postal code: Z00P5Y4

Address: Kazakhstan, Nur-Sultan, Abylai Khan Avenue, 15A

Phone: +77479894746

E-mail: oktyabrovdana@gmail.com

J Trauma Ortho Kaz 2021; 3 (59): 4-8

Recieved: 24-07-2021

Accepted: 15-08-2021



This work is licensed under a Creative Commons Attribution 4.0 International License

Introduction

Developmental dysplasia of the hip (DDH) occurs among 2-3 in 1000 of infants. DDH influences to the biomechanics of the hip and leads to early development of coxarthrosis [1,2]. Dysplastic coxarthrosis among adults develops in 21-80% rate in case of congenital pathology of the hip and is the most common surgery pathology among kids and teenagers, that occurs, according to different resources, at 0,7-2,5% case in 1000 of infants [3-5]. According to Guo C.Y. et al., (2015), approximately 40% of dysplastic coxarthrosis of the hip among adults is the consequence of joint development defects untreated in childhood [6].

Dysplastic coxarthrosis is the basic reason of total hip arthroplasty (THA) among young people (from 21% to 29%) [7].

The degree of femur displacement is specified by different classifications Crowe J.F. (1979) [8], Hartofilakidis G (1988) [9], Eftekhari N.S. (1978) [10]. In these classifications, the final stages of the displacement of the femoral head corresponds to a developmental displacement of the hip. Consequently G. Hartofilakidis at his work in 2004, describes two subcategories at types of high dislocation to clarify the relationship between the femoral head and the formed false trough on the wing of the iliac bone – supporting (Crowe IV, Hartofilakidis C1) or unsupported (Crowe IV, Hartofilakidis C2).

Medical-social side of the question, connected with the life quality of given patients group is also the major problem for healthcare systems of different countries in

the world. Each year, the number of patients with dysplastic coxarthrosis who require expensive THA with further long-term rehabilitation is growing gradually [12,13]. Zhaksybaev M.H. et al., (2016) noted that in Kazakhstan, as well as in the rest of the world, the number of surgery on replacement of major joints is increasing. There is a necessity of 24500 surgeries of THA on hip and knee joints annually in the country [14].

The features of THA at dysplastic coxarthrosis depends on the degree of dysplasia and is followed by the possibility of several issues development. However, at the dysplastic coxarthrosis, as a rule, THA gives worse results of curing, as well as at post-traumatic coxarthrosis. Surgery in this case is more complex and the and disruption of normal anatomy leads to an increased incidence of surgical errors [15-17]. According to literature, in 10% cases, the THA can lead to periprosthetic infections, deep vein thrombophlebitis of the operated limb. In the postoperative period, dislocation of the head of the endoprosthesis can take place, instability of prosthesis, which subsequently requires re-arthroplasty.

Taking into consideration the frequency and impact on the quality of life of patients of working age, dysplastic coxarthrosis is indeed relevant issue for modern orthopedics and is required to be studied in Kazakhstan.

The aim of the review – to research the prevalence and medical-social aspects of dysplastic coxarthrosis as consequence of developmental dysplasia of the hip in the world and Kazakhstan.

The prevalence of dysplastic coxarthrosis

The frequency of coxarthrosis in the world is distributed unequally. As a result of Randall T. Loder et al. (2011) systematic review, the prevalence of coxarthrosis is different: In Europe, 7-25% of people older than 55 years are suffering from coxarthrosis at different etiology, as well these indicators as lower for Asian and Afro-American [19].

This opinion is justified by results of our literature search. Based on analyzed literature resources, South Korea showed the lowest results of coxarthrosis (78,5⁰/₀₀₀₀) in last ten years. The highest results are in Switzerland - 356,5⁰/₀₀₀₀, Germany - 299,1⁰/₀₀₀₀, France - 277,7⁰/₀₀₀₀, UK - 214,4⁰/₀₀₀₀, Poland - 142⁰/₀₀₀₀, Turkey - 99,4⁰/₀₀₀₀. In Russia, this is resulted at 17.8-20.0 cases in 10 000 adults [20-22].

The DDH cases among racial groups also differs based on geographical location. It is noted, that tight swaddling of infants has a non-balgoric effect on the condition of the hip joint [19].

According to foreign literature, dysplastic coxarthrosis takes from 25% to 77% at the structure of degenerative-dystrophic diseases of the hip joint [23-25]. In average, the dysplastic affection of the hip accounts for 16.5% of all pathology of the organs of support and motion [25].

Medical-social questions on dysplastic coxarthrosis

Progressive feature of hip joint destruction is leading to decreased working abilities in 60% and to disablement in 11,5% [27,28]. We decided to research the medical-social side of the question from perspective of non-surgical and patient that went through arthroplasty.

Concluding, the coxarthrosis of dysplastic genesis is accounting for 50% of osteoarthritis of large joints. The prevalence of pathology and the clinical severity at II

According to Batpen A. et al., (2013), the prevalence of coxarthrosis in Kazakhstan is also unequally distributed. The lowest coefficient is in Atyrau state (1,7%), the highest in Mangystau state (47,0%) and Almaty state (40,9%) [26]. The found unequal distribution of rates is connected with the level of healthcare support, specifically with effective registration and patient screening and later finding DDH. Unfortunately, the distribution of dysplastic form of coxarthrosis in Kazakhstan is not yet researched in detail.

According to conducted analysis of patients with dysplastic coxarthrosis through National Scientific Center of Traumatology and Orthopaedics named after Academician Batpenov N.D. (Former name - Scientific Center of Traumatology and Orthopaedics) in the period of 2015-2020, the distribution of dysplastic coxarthrosis is from 25% to 30% in the structure of degenerative-dystrophic illnesses of the hip among adult population of Kazakhstan. In Crowe classification, I-II type - 72,2%, III-IV type - 27,8%.

and III stages of dysplastic coxarthrosis defines the high medical-social important of its prevention and treatment. The developmental inferiority of the hip joint tissues and overloading of the joint components due to a deficiency in coverage of the femoral head by the acetabulum play major at pathogenesis of dysplastic coxarthrosis [29].

The severance and disability occurred by DDH and hip joints is influenced by late diagnosis and starting

point of treatment, unjustified long treatment through conservative methods, application of repeated closed femoral head repositioning. As a result of numerous corrections of femoral head, the deformation and dystrophia of femoral head is developed. The late DDH process is also defined by palliative characteristics, and it improves the ability to support the limb for a short period of time [29].

According to the literature, the group of disabled children with Congenital dislocation of the hip is dominated by girls (1:3). The average age of disabled children affected by dysplastic coxarthrosis is 11,6 years. Among adult population, there is a domination of women (1:4,3). The average age of disabled people with dysplastic coxarthrosis is 40 years. The average weight of disabled after congenital dislocation of the hip that undertook conservative treatments is higher (70,1%) than those after THA [27,28].

Results of life quality evaluation of patients with dysplastic coxarthrosis is decreasing as the child gets older.

Conclusions

Dysplastic coxarthrosis incidence rate in the world differs depending on the geographical and racial groups, while it is lower among males than females. The number of dysplastic coxarthrosis patients that are waiting for full arthroplasty with further long-term rehabilitation is increasing gradually.

It is connected with the increase in weight, deformation, and decreased functionality of hip joints from one or both sides [29]. After conducting THA, the life quality increases dramatically. It was widely believed that the effective tool for measuring the life quality of such group was the scale and test of Chanley and Harris et al., that are focused on clinical-functional condition of surgical joint. However, several authors believe that given scale and tests are not also effective tool for concluding the impact of treatment on the quality of life [29-31]. According to our evaluation results, the quality of life after THA is indeed higher than those non-surgical. However, it is important to note that the quality of life that undergo THA is directly depending on the effective and relevant rehabilitation [32,33]. Because of this, providing accessible rehabilitation actions is still an important question at most of developing countries, as well as Kazakhstan.

There is a necessity in research of the prevalence, as well as conducting the medical-social evaluation of the consequences among those with dysplastic coxarthrosis in Kazakhstan.

References

1. Vaquero-Picado A., González-Morán G., Garay E.G., Moraleta L. *Developmental dysplasia of the hip: update of management*. *EFORT open reviews*. 2019; 4(9): 548-556. <https://doi.org/10.1302/2058-5241.4.180019>.
2. Shaw B.A., Segal L.S. *Evaluation and Referral for Developmental Dysplasia of the Hip in Infants*. *Pediatrics*. 2016; 138(6): e20163107. <https://doi.org/10.1542/peds.2016-3107>.
3. Harsanyi S., Zamborsky R., Krajciová L., Kokavec M. et al. *Developmental dysplasia of the hip: a review of etiopathogenesis, risk factors, and genetic aspects*. *Medicina*. 2020; 56(4): 153. <https://doi.org/10.3390/medicina56040153>.
4. Eftekhari N.S. *Congenital dysplasia and dislocation in total hip arthroplasty*. St. Louis: Mosby. 1993. P. 925-963.
5. Bohaček I., Plečko M., Duvančić T., Smoljanović T. et al. *Current knowledge on the genetic background of developmental dysplasia of the hip and the histomorphological status of the cartilage*. *Croatian Medical Journal*. 2020; 61(3): 260-270. <https://doi.org/10.3325/cmj.2020.61.260>.
6. Guo C.Y., Liang B.W., Sha M., Kang L.Q. et al. *Cementless arthroplasty with a distal femoral shortening for the treatment of Crowe type IV developmental hip dysplasia*. *Indian journal of orthopaedics*. 2015; 49(4): 442-446. <https://doi.org/10.4103/0019-5413.159652>.
7. Gkiatas I., Boptsi A., Tserga, D., Gelalis I. et al. *Developmental dysplasia of the hip: a systematic literature review of the genes related with its occurrence*. *EFORT open reviews*, 2019; 4(10): 595-601. <https://doi.org/10.1302/2058-5241.4.190006>.
8. Crowe J.F., Mani V.J., Ranawat C.S. *Total hip replacement in congenital dislocation and dysplasia of the hip*. *J Bone Joint Surg Am*. 1979; 6(1): 15-23.
9. Hartofilakidis G., Stamos K., Ioannidis T.T. *Low friction arthroplasty for old untreated congenital dislocation of the hip*. *J Bone Joint Surg Br*. 1988; 70(2): 182-186. <https://doi.org/10.1302/0301-620X.70B2.3346284>.
10. Eftekhari N.S. *Congenital dysplasia and dislocation in total hip arthroplasty*. St. Louis: Mosby. 1993. P. 925-963.
11. Hartofilakidis G., Karachalios T. *Total hip arthroplasty for congenital hip disease*. *JBJS*. 2004; 86(2): 242-250. <https://doi.org/10.2106/00004623-200402000-00005>.
12. Kawasaki M., Hasegawa Y., Okura T., Ochiai S. et al. *Muscle damage after total hip arthroplasty through the direct anterior approach for developmental dysplasia of the hip*. *J Arthroplasty*. 2017; 32(8): 2466-2473. <https://doi.org/10.1016/j.arth.2017.03.043>.
13. Шнайдер Л.С., Голеньков О.И., Тургунов Э.У., Ефименко М.В. и др. *Укорачивающая подвертельная остеотомия бедренной кости при эндопротезировании тазобедренного сустава у пациентов с врожденным вывихом бедра // Генный ортопедии. – 2020. – Т. 26. – №3. – С. 340-346.* <https://doi.org/10.18019/1028-4427-2020-26-3-340-346>.
- Shnaider L.S., Golenkov O.I., Turgunov E.U., Efimenko M.V. i dr. *Ukorachivaiushchaia podvertel'naia osteotomiia bedrennoi kosti pri endoprotezirovanii tazobedrennogo sustava u patsientov s vrozhdenным vyvikhom bedra (Shortening collapsed osteotomy of the femur for hip arthroplasty in patients with congenital hip dislocation) [in Russian]. *Genii ortopedii*. 2020; 26(3): 340-346. <https://doi.org/10.18019/1028-4427-2020-26-3-340-346>.*
14. Джаксыбаев М.Н., Оразхан Ж., Нурлыбекова Е.Н., Ушуров А.А. и др. *Анализ эндопротезирования по поводу дисплазии тазобедренного сустава // Вестник Казахского национального медицинского университета. – 2016. – №3-1. – С. 245-249.*
- Dzhakysybaev M.N., Orazkhan Zh., Nurlybekova E.N., Ushurov A.A. i dr. *Analiz endoprotezirovaniia po povodu displazii tazobedrennogo sustava (Analysis of arthroplasty for hip dysplasia) [in Russian]. *Vestnik Kazakhskogo natsional'nogo meditsinskogo universiteta*. 2016; 3-1: 245-249.*

15. Волокитина Е.А. Эндопротезирование тазобедренного сустава после опорных остеотомий бедренной кости // Современное искусство медицины. – 2013. – №6(14). – С. 10–15.
 Volokitina E.A. Endoprotezirovanie tazobedrennogo sustava posle opornykh osteotomii bedrennoi kosti (Endoprosthetics of the hip joint after supporting osteotomies of the femur) [in Russian]. *Sovremennoe iskusstvo meditsiny*. 2013; 6(14): 10–15.
16. Гольник В.Н., Григоричева Л.Г., Джухаев Д.А., Меркулов С.А. и др. Реконструкция вертлужной впадины при сложных случаях первичного эндопротезирования тазобедренного сустава // Основные направления отечественной травматологии и ортопедии: Материалы Крымского форума травматологов-ортопедов. – 2016. – С. 137–140.
 Gol'nik V.N., Grigoricheva L.G., Dzhukhaev D.A., Merkulov S.A. i dr. Rekonstruktsiia vertluzhnoi vpadiny pri slozhnykh sluchaiakh pervichnogo endoprotezirovaniia tazobedrennogo sustava (Reconstruction of the acetabulum in complex cases of primary hip arthroplasty) [in Russian]. *Osnovnye napravleniia otechestvennoi travmatologii i ortopedii: Materialy Krymskogo foruma travmatologov-ortopedov*. 2016; 137–140.
17. Negrin L.L., Seligson D. Results of 167 consecutive cases of acetabular fractures using the Kocher-Langenbeck approach: a case series. *J Orthop Surg Res*. 2017; 12(1): 66. <https://doi.org/10.1186/s13018-017-0563-6>.
18. Решетников А.Н., Павленко Н.Н., Зайцев В.А., Фроленков А.В. и др. Тотальное эндопротезирование тазобедренного сустава при диспластическом коксартрозе // Вестник Тамбовского университета. Серия: естественные и технические науки. – 2012. – Т. 17. – №3. – С. 901–603.
 Reshetnikov A.N., Pavlenko N.N., Zaitsev V.A., Frolenkov A.V. i dr. Total'noe endoprotezirovanie tazobedrennogo sustava pri displasticheskom koksartroze (Total hip arthroplasty for dysplastic coxarthrosis) [in Russian]. *Vestnik Tambovskogo universiteta. Serii: estestvennye i tekhnicheskie nauki*. 2012; 17(3): 901–603.
19. Loder R.T., Skopelja E.N. The epidemiology and demographics of hip dysplasia. *International Scholarly Research Notices*. 2011; 2011: 238607. <https://doi.org/10.5402/2011/238607>.
20. Swain S., Sarmanova A., Mallen C., Kuo C.F. et al. Trends in incidence and prevalence of osteoarthritis in the United Kingdom: findings from the Clinical Practice Research Datalink (CPRD). *Osteoarthritis and cartilage*. 2020; 28(6): 792–801. <https://doi.org/10.1016/j.joca.2020.03.004>.
21. Saltychev M., Perna K., Seppänen M., Mäkelä K. et al. Pelvic incidence and hip disorders: A systematic review and quantitative analysis. *Acta orthopaedica*. 2018; 89(1): 66–70. <https://doi.org/10.1080/17453674.2017.1377017>.
22. Королько А.С. Суцевич В.В., Евко Я.И., Кезля О.П. Популяционная характеристика пациентов с коксартрозом // Известия Национальной академии наук Беларуси. Серия медицинских наук. – 2019. – Т. 16. – №2. – С. 156–165. <https://doi.org/10.29235/1814-6023-2019-16-2-156-165>.
- Korol'ko A.S., Sushchevich V.V., Evko Ia.I., Kezlia O.P. Populiatsonnaia kharakteristika patsientov s koksartrozom (Population characteristics of patients with coxarthrosis) [in Russian]. *Izvestiia Natsional'noi akademii nauk Belarusi. Serii meditsinskikh nauk*. 2019; 16(2): 156–165. <https://doi.org/10.29235/1814-6023-2019-16-2-156-165.22>.
23. Корьяк В.А., Сороковиков В.А., Свистунов В.В., Шарова Т.В. Эпидемиология коксартроза // Сибирский медицинский журнал (Иркутск). – 2013. – Т. 123. – №8. – С. 39–44.
 Kor'iak V.A., Sorokovikov V.A., Svistunov V.V., Sharova T.V. Epidemiologiia koksartroza (Epidemiology of coxarthrosis) [in Russian]. *Sibirskii meditsinskii zhurnal (Irkutsk)*. 2013; 123(8): 39–44.
24. Сергеев С.В., Жмотова Е.А., Киммельфельд И.М., Золотухина И.Д. и др. Эволюция коксартроза в свете экспертизы трудоспособности // Вестник травматологии и ортопедии им. Н.Н. Приорова. – 1996. – Т. 3. – №2. – С. 3–10.
 Sergeev S.V., Zhmotova E.A., Kimmel'fel'd I.M., Zolotukhina I.D. i dr. Evoliutsiia koksartroza v svete ekspertizy trudospobnosti (The evolution of coxarthrosis in the light of the examination of the ability to work) [in Russian]. *Vestnik travmatologii i ortopedii im. N.N. Priorova*. 1996; 3(2): 3–10.
25. Eskelinen A. Total hip arthroplasty in young patients - with special references to patients under 55 years of age and to patients with developmental dysplasia of the hip. Helsinki. 2006; 128.
26. Brunner B., Ulmar H., Reichel R., Decking R. Decking The Eftekhar and Kerboul classification in assessment of developmental dysplasia of the hip in adult patients. Measurement of inter- and intraobserver reliability. *HSS Journal: the Musculoskeletal Journal of Hospital for Special Surgery*. 2007; 4(1): 25–31. <https://doi.org/10.1007/s11420-007-9066-z>.
27. Igissinov N., Baimagambetov S., Batpen A. Evolutionary epidemiological assessment of coxarthrosis incidence among adults population in Kazakhstan. *World Appl Sci J*. 2013; 24(9): 1271–1275. <https://doi.org/10.5829/idosi.wasj.2013.24.09.13276>.
28. Savchenko V., Maykova T., Afanasiev S., Kashuba V. et al. Disorders of the mineral exchange and metabolism of bone tissue as a pathogenetic basis of physical rehabilitation patients with coxartrosis. *Journal of Physical Education and Sport*. 2020; 20(1): 447–451. <https://doi.org/10.7752/jpes.2020.s1065>.
29. Dogaru G. The importance of the elemental functional mobility coefficient in assessing the functional status of the coxofemoral joint. *Balneo Research Journal*. 2018; 9(1): 38–42. <https://doi.org/10.12680/balneo.2018.169>.
30. Akberdina D.L. Etiology and pathogenesis of coxarthrosis and methods of its treatment. *Kazan medical journal*. 1983; 64(6): 404–408. <https://doi.org/10.17816/kazmj83602>.
31. Świtoń A., Wodka-Natkaniec E., Niedźwiedzki Ł., Gaździk T. et al. Activity and Quality of Life after Total Hip Arthroplasty. *Ortopedia, traumatologia, rehabilitacja*. 2017; 19(5): 441–450. <https://doi.org/10.5604/01.3001.0010.5823>.
32. Balik M.S., Hocaoglu Ç., Erkut A., Güvercin Y. et al. Evaluation of the quality of life and psychiatric symptoms of patients with primary coxarthrosis after total hip arthroplasty. *Acta Chir Orthop Traumatol Cech*. 2017; 84(6): 436–40.
33. Kraydjikova L., Nikolovska L., Krstev T., Stratovska T. Physiotherapy program for improving the quality of life in patients with coxarthroses. *Journal of Biomedical and Clinical Research*. 2015; 8(1): 69–70.

Диспластикалық коксартроздың эпидемиологиялық және медициналық-әлеуметтік аспектілері

Октяброва Д.Г. ¹, Әшімов Қ.Д. ², Байдалин Т.Т. ³, Сүлейменов Б.Т. ⁴, Бекназаров А.И. ⁵

¹ №3 ортопедия бөлімшесінің дәрігер резиденті, Академик Батпенев Н.Д. атындағы Травматология және ортопедия ұлттық ғылыми орталығы, Астана медицина университетінің PhD докторанты, Нұр-Сұлтан, Қазақстан.

E-mail: oktyabrovanadana@gmail.com

² №3 ортопедия бөлімшесінің меңгерушісі, Академик Батпенев Н.Д. атындағы Травматология және ортопедия ұлттық ғылыми орталығы, Нұр-Сұлтан, Қазақстан. E-mail: kairat_ashimov@mail.ru

³ №7 ортопедия бөлімшесінің меңгерушісі, Академик Батпенев Н.Д. атындағы Травматология және ортопедия ұлттық ғылыми орталығы, Нұр-Сұлтан, Қазақстан. E-mail: t.baidalin@gmail.com

⁴ №3 ортопедия бөлімшесінің дәрігер травматолог-ортопеді, Академик Батпенев Н.Д. атындағы Травматология және ортопедия ұлттық ғылыми орталығы, Нұр-Сұлтан, Қазақстан. E-mail: bekjan-suleimen@mail.ru

⁵ №3 ортопедия бөлімшесінің дәрігер травматолог-ортопеді, Академик Батпенев Н.Д. атындағы Травматология және ортопедия ұлттық ғылыми орталығы, Нұр-Сұлтан, Қазақстан. E-mail: askar_big_tan@mail.ru

Түйіндеме

Әлемдегі диспластикалық коксартрозбен аурушаңдық жиілігі географиялық орналасуына және нәсілдік топқа байланысты айтарлықтай өзгереді. Ерлер мен әйелдердегі аурушаңдық көрсеткіші шамамен бірдей деңгейде екені байқалады. Диспластикалық коксартрозбен ауыратын науқастардың ішіндегі әрі қарай ұзақ мерзімді оңалтуды қажет ететін толық эндопротездеуді күтетіндердің саны тұрақты өсіп келеді.

Қазақстанда диспластикалық коксартрозбен ауыратын науқастардың таралуын егжей-тегжейлі зерттеу, сондай-ақ, аурудың салдарын медициналық-әлеуметтік бағалау қажет.

Түйін сөздер: жамбастың туа біткен дисплазиясы, диспластикалық коксартроз, өмір сапасы, мүгедектік, жас науқастар.

Эпидемиологические и медико-социальные аспекты диспластического коксартроза

Октяброва Д.Г. ¹, Ашимов К.Д. ², Байдалин Т.Т. ³, Сулейменов Б.Т. ⁴, Бекназаров А.И. ⁵

¹ Врач резидент отделения ортопедии №3, Национальный научный центр травматологии и ортопедии имени академика Батпенева Н.Д., PhD докторант Медицинского университета Астана, Нур-Султан, Казахстан.

E-mail: oktyabrovanadana@gmail.com

² Заведующий отделением ортопедии №3, Национальный научный центр травматологии и ортопедии имени академика Батпенева Н.Д., Нур-Султан, Казахстан. E-mail: kairat_ashimov@mail.ru

³ Заведующий отделением ортопедии №7, Национальный научный центр травматологии и ортопедии имени академика Батпенева Н.Д., Нур-Султан, Казахстан. E-mail: t.baidalin@gmail.com

⁴ Врач травматолог-ортопед отделения ортопедии №3, Национальный научный центр травматологии и ортопедии имени академика Батпенева Н.Д., Нур-Султан, Казахстан. E-mail: bekjan-suleimen@mail.ru

⁵ Врач травматолог-ортопед отделения ортопедии №3, Национальный научный центр травматологии и ортопедии имени академика Батпенева Н.Д., Нур-Султан, Казахстан. E-mail: askar_big_tan@mail.ru

Резюме

Показатели заболеваемости диспластическим коксартрозом в мире значительно различаются в зависимости от географического положения и расовой группы. Показатели заболеваемости у мужчин и женщин примерно на одном уровне. Количество пациентов с диспластическим коксартрозом, ожидающих проведения тотального эндопротезирования с дальнейшей длительной реабилитацией неуклонно растет.

Есть необходимость детального изучения распространенности, а также проведения медико-социальной оценки последствий заболевания у больных диспластическим коксартрозом в Казахстане.

Ключевые слова: врожденная дисплазия бедра, диспластический коксартроз, качество жизни, инвалидность, молодой возраст.